


I-Valex[®]-1

Amino acid-modified infant formula with iron

Nutrition support of infants and toddlers with a disorder of leucine catabolism. Use under medical supervision.



Features

- Leucine-free to allow greater intake of intact protein.
- Isoleucine and valine to help prevent deficiency of these amino acids.
- L-carnitine (900 mg/100 g) to help supply amount normally found in human milk and foods of animal origin and to help excrete toxic metabolites.
- Glycine (1000 mg/100 g) to bind with acyl groups and help enhance their excretion in a nontoxic form.
- Approximately 40% of energy as fat to help achieve acceptable formula osmolality.
- 6.6% of energy as linoleic acid.
- Nutrient profile specifically designed for infants and toddlers.
- When fed according to the *Abbott Nutrition Support Protocols*, provides adequate amounts of all nutrients.
- Non-GMO.*
- DHA and ARA to help support brain and eye development.
- Lactose-free.
- Gluten-free.
- Halal.

* Ingredients not genetically engineered.

Safety Precautions

- Not intended as a sole source of nutrition.
- Give only to infants and toddlers with proven disorders of leucine catabolism who are under medical supervision.
- Must be supplemented with protein and fluid in prescribed amounts to completely meet leucine and water requirements.
- Do not boil mixture or use terminal sterilization.
- **Never use a microwave oven to warm formula.** Serious burns can result.
- Not for IV use.
- Powdered infant formulas are not sterile and should not be fed to premature infants or infants who might have immune problems unless directed and supervised by your baby's doctor.

Availability

List Number	Item
67044	I-Valex-1 / 14.1 oz (400 g) Can / 6 ct

Ingredients

Unflavored Powder

Corn Syrup Solids, High Oleic Safflower Oil, Coconut Oil, Soy Oil, L-Alanine, L-Glutamic Acid, Modified Corn Starch. Less than 2% of: L-Aspartic Acid, L-Proline, L-Lysine, Calcium Phosphate, L-Arginine, Potassium Phosphate, Glycine, L-Carnitine, L-Tyrosine, L-Phenylalanine, L-Serine, L-Threonine, Sodium Citrate, Potassium Citrate, L-Valine, Magnesium Chloride, L-Isoleucine, L-Histidine, L-Methionine, M. Alpina Oil, C. Cohnii Oil, Calcium Carbonate, Citric Acid, Ascorbic Acid, L-Cystine Dihydrochloride, L-Tryptophan, Inositol, Choline Chloride, Taurine, Ferrous Sulfate, Ascorbyl Palmitate, Zinc Sulfate, dl-Alpha-Tocopheryl Acetate, Mixed

Tocopherols, Calcium Pantothenate, Niacinamide, Manganese Sulfate, Thiamine Hydrochloride, Copper Sulfate, Vitamin A Palmitate, Riboflavin, Pyridoxine Hydrochloride, Potassium Iodide, Folic Acid, Phylloquinone, Chromium Chloride, Sodium Selenate, Sodium Molybdate, Biotin, Vitamin D3, Vitamin B12, and Salt.

Nutrition Information

	100 g Powder	
	Value	%DV
Protein Equivalent, g	15.0	
Fat, g	21.7	
Carbohydrate, g	53.0	
Linoleic Acid, mg	3650	
Linolenic Acid, mg	360	
L-Carnitine, mg	900	
Calories	480	
Vitamin A, IU	1300	
Vitamin A, mcg RE	390	
Vitamin D, IU	450	
Vitamin D, mcg	11.3	
Vitamin E, IU	12.5	
Vitamin E, mg alpha-TE	6.4	
Vitamin K, mcg	60	
Thiamin (Vitamin B1), mcg	1120	
Riboflavin (Vitamin B2), mcg	650	
Vitamin B6, mcg	560	
Vitamin B12, mcg	0.85	
Niacin, mcg	4450	
Niacin, mg NE	7.7	
Folic Acid (Folacin), mcg	88	
Pantothenic Acid, mcg	4000	
Biotin, mcg	23.5	
Vitamin C, mg	75	
Choline, mg	95	
Inositol, mg	144	
Calcium, mg	575	
Phosphorus, mg	400	
Magnesium, mg	50	
Iron, mg	7.6	
Zinc, mg	5.2	

Manganese, mcg	385
Copper, mcg	408
Iodine, mcg	65
Selenium, mcg	21
Chromium, mcg	12
Molybdenum, mcg	13.8
Sodium, mg	190
Sodium, mEq	8.3
Potassium, mg	675
Potassium, mEq	17.3
Chloride, mg	325
Chloride, mEq	9.2

Preparation

Instructions for Use: Use Only As Directed by a Physician

Your baby's health depends on carefully following these directions. Proper hygiene, handling and storage are important when preparing infant formula. Failure to follow these directions could result in severe harm. Ask your baby's doctor if you need to use cooled, boiled water for mixing and if you need to boil (sterilize) bottles, nipples and rings before use.

- Wash your hands, surfaces and utensils.
- Pour prescribed amount of water into clean container.
- Add prescribed amount of I-Valex-1 powder (and other ingredients if recommended).
- Shake well for 10-15 seconds; if using a blender, mix no more than 5 seconds.
- Pour mixture into clean feeding bottles or container; cap.
- Once feeding begins, use within 1 hour or discard.

Approximate weights for unpacked, level U.S. standard dry household measures for powder*

Household Measure (US)	Weight (g)
1 Tbsp	8
1/4 Cup	30
1/3 Cup	40
1/2 Cup	60
1 Cup	120

* For most accurate results, I-Valex-1 should be weighed on a scale that reads in grams.

Storage & Handling

Instructions for Use:

- Once mixed, store bottles or container in refrigerator and **feed to baby or child within 24 hours**
- Store unopened or opened can at room temperature; avoid extreme temperatures.
- **Use opened can contents within 1 month.**