

A hypoallergenic amino acid-based formula with a unique lipid blend for infants with cow's milk protein allergy (CMPA), multiple food allergies, or malabsorptive conditions



# Alfamino

# 0-12 months of age

## DIRECTIONS FOR PREPARATION AND USE:

Always wash your hands and utensils before preparing formula. For oral or tube feeding. Use under medical supervision.



STEP 1 Pour desired amount of water into bottle. See chart below.



STEP 2 Add 1 unpacked level scoop (4.7g) of powder for each 1 fl oz water. (Only use enclosed scoop.)



STEP 3 Cap bottle; SHAKE WELL. Feed baby immediately. Once feeding begins, use contents within 1 hour or discard.

# MIXING GUIDE:

TO MAKE APPROX.:	MEASURE WATER	ADD POWDER	
<b>2</b> fl oz bottle	<b>2</b> fl oz	<b>2</b> unpacked level scoops	<b>NOTE:</b> Only if necessary, refrigerate freshly prepared bottle and use contents within 24 hours. Refrigerator temperatures should be maintained between 35-40° F (2-4° C). Re-shake bottle before feeding.
<b>4</b> fl oz bottle	<b>4</b> fl oz	4 unpacked level scoops	
<b>6</b> fl oz bottle	<b>6</b> fl oz	<b>6</b> unpacked level scoops	
<b>8</b> fl oz bottle	<b>8</b> fl oz	8 unpacked level scoops	

**SCOOP WEIGHT:** 4.7g. 1 unpacked level scoop added to 1 fl oz of water makes approximately 1 fl oz. When mixed as directed, makes approximately 94 fl oz (2.9qt)

## WARNING: Your baby's health depends on carefully following the preparation, use, and storage instructions. Failure to follow these instructions could result in severe harm.

Consult your baby's doctor about the appropriate formula for your baby. Powdered infant formulas are not sterile and should not be fed to premature infants or infants who may have immune problems unless directed and supervised by your baby's doctor. Sterilization or boiling of tap or bottled water and sterilization of utensils is recommended. Your doctor will advise if it is not required. If boiling water, bring to a rolling boil for 1 minute and cool before preparing formula.

#### Do not use microwave to warm formula. Serious burns may result.

**STORAGE:** Tightly cover open can and use within 1 month. Store open and unopened cans in a cool, dry place. Avoid extreme storage temperatures.

IMPORTANT NOTICE: Mothers should be encouraged to continue breastfeeding even when their infants have cow's milk protein allergy. This usually requires dietary counseling to completely exclude all sources of cows' milk protein from the mother's diet. If a decision to use specialty formulas is taken, it must be used under medical supervision.

